MICROWAVE BIOEFFECT CONGRUENCE WITH SCHIZOPHRENIA

John J. McMurtrey,^a Copyright 2002, 8/25/03

ABSTRACT

In view of the documentation for microwave technology to induce internal voice in humans, the correlation between microwave bioeffects and schizophrenia is reviewed. These correlations are extensive and include cognitive deficit, electrophysiologic activity, startle decrease, neurotransmitter changes, hormone alterations, immune alterations, mitochondria deficits, deleterious histologic change in disease reduced brain areas, activation of hallucination involved brain areas, and ocular disease. Schizophrenia correlates with microwave bioeffests such that congruence indicates microwave involvement with this disorder. The development of methods to exclude microwave means in psychosis is imperative, and research is proposed.

INTRODUCTION

Microwave induced sound¹and internal voice technology has long been discovered, developed,^{2, 3} detailed in patents, ^{,4} with weapons applications described. ^{,5, 6} That such technology can be applied remotely and coupled to target tracking technology⁷ has implications for patients who, by virtue of internal voice complaint and other symptoms are diagnosed with schizophrenia.⁸ A frequent patient understanding of the origin of these voices is by remote transmission, though the very concept is considered delusional,⁹ and

^a Address: 903 N. Calvert St., Baltimore MD 21202. Email- <u>Johnmcmurt@aol.com</u> Phone- 410-539-5140.

often the diagnosis is psychosis of varying severity depending on functional ability,¹⁰ without any investigation of the described internal voice capabilities.

Microwave voice transmission substantiation suggests examination of microwave bioeffect correlation with schizophrenia, which has the most hallucination prevalence. This examination reveals correlations that are listed in Table I. These correlations are so numerous and extensive that a high level of congruence between microwave bioeffects and schizophrenia is apparent. The effects discussed are within the microwave hearing spectrum, (100-10,000 MHz or 0.1-10 GHz) and intensity is in terms of the existing US standards,¹¹ as in parenthesis. Microwave exposure duration terms are defined below.^b

Cognitive Deficit

Schizophrenia cognitive deficit is particularly in memory.^{12, 13} Schizophrenic prefrontal cortex dysfunction is considered important, with this region's 'working memory' role as central, to many schizophrenic symptoms.¹⁴ Rat "working memory" performance in a radial arm maze is deficient on microwave exposure (60% of US pop. std.) ^{15, 16} Rat water maze performance is deficient in 'spatial memory' with acute pulsed microwave exposure (1.2 X US pop. std.)¹⁷ Prolonged rat microwave continuous wave exposures exhibit shuttle box and intermittent response training deficits (30% of US pop. std. to 1.2 X US occup. std.),^{18, 19, 20, 21} with more pronounced decrements on extended exposure.²² Other studies found rat food pellet reinforcement deficits on continuous wave or pulsed microwave exposures (½ to 1.6 X US occup. std.)^{23, 24, 25} Multiple human case report of memory difficulty, with other neurasthenic complaints exists on excess microwave exposure.²⁶

^b Brief = 30 minutes or less; Acute = 60 minutes to 30 minutes; Prolonged or Extended = days, Chronic = month or more.

Accidental and/or occupational 1-10 GHz excess radar exposure exhibits frontal lobe neuropsychiatric symptoms.²⁷

Electrophysiology

An electrophysiologic indicator of 'working memory', contingent negative variation (CNV)²⁸ is decreased in several mental illnesses with the greatest decreases in schizophrenia.^{29, 30, 31} Cell phone radiation also decreases human CNV.^{32, 33} The test involves a warning stimulus and an imperative stimulus with the intervening evoked waveform representative of sensory and motor adjustment prior to expected action.

Electrophysiologic auditory event related P300 and antecedents are reduced in schizophrenia,^{34, 35} with increased latency.³⁶ Magnetoencephalographic (MEG) auditory event responses during hallucination are also decreased,³⁷ resembling responses to interfering sound.³⁸ Like hallucination or outside sound, pulsed microwaves decrease auditory electrophysiologic evoked potential amplitudes with a tendency towards increased latency in rats and rabbits (less than US occup. std.)³⁹ Hearing effect pulsed microwaves evoke brain responses similar to auditory stimuli.^{40, 41, 42} Human hearing threshold increases for auditory tones on radiofrequency exposure.⁴³ Schizophrenia auditory P300 reduction is related to deleterious signs and poor prognosis.⁴⁴

Soviet and American microwave exposure of humans report EEG increases in delta or "slow" waves, abnormal to alertness in quantity. Acute human exposure to continuous waves at the low microwave hearing spectrum end and pulsed frequencies at the high spectrum end, exhibit increased electroencephalogram (EEG) delta waves (less than US pop. std.)⁴⁵ Soviet and East European microwave occupational exposure review observes increased EEG delta waves.⁴⁶ Cell phones also increase human delta waves.⁴⁷ Rabbit and rat microwave irradiation yield delta waves as well. Daily 3 hour rabbit exposures produces delta wave increases at 1 month to pulsed microwaves and at 2 months to continuous wave exposure (1/2 US occup. std.)⁴⁸ Daily 7 hour microwaves produces delta waves after 10-15 days in rabbits at 1/3rd the US population exposure standard, but took 1 month for delta wave increase at 1/30th this standard.⁴⁹ Rat microwave irradiation induces delta waves in the left hemisphere by continuous wave, but in the right hemisphere when modulated.⁵⁰ Delta waves are also produced by extra low frequency radiation in rabbits⁵¹ or magnetic fields in humans.⁵²

Microwave delta wave increases correspond to delta wave increases widely noted in untreated, ^{53, 54, 55, 56, 57, 58, 59, 60, 61} and medicated^{62, 63, 64, 65, 66, 67, 68, 69} schizophrenia EEGs.⁷⁰ Delta waves particularly correspond to psychotic episodes, ^{71, 72} and occur immediately prior to auditory hallucination.⁷³

Electromagnetic field EEG entrainment occurs especially within physiologic brain frequencies (1-40 Hz.), either with a so modulated carrier wave or at these extra low frequencies. Microwave EEG entrainment (or change to exposure frequency) is demonstrated in cats,⁷⁴ and rats.⁷⁵ Lower frequency radiation or magnetic EEG entrainment is observed in rabbits,⁷⁶ monkeys,⁷⁷ and humans.⁷⁸ In addition to the capacity of entrainment to produce delta waves, the effect forms a basis for schizophrenic thought interference complaints, and is of non-lethal weapon concern.⁷⁹

Startle Response

Some schizophrenics are hypo- or non-responders to orienting responses⁸⁰ and normally evoked electrodermal activity.⁸¹ Microwave occupational exposure inhibits galvanic skin response.⁸² Some schizophrenics have little or no startle response.⁸³ Microwave exposed rats exhibit decreased startle under both continuous wave⁸⁴ and pulsed^{85, 86} conditions (1.2 X US occup. std.) Pre-natal rat exposure decreases startle in females (1.2 X US occup. std.)⁸⁷ Rats also fight less on microwave exposure (23 % of US pop. std.),⁸⁸ and avoid hearing effect pulsed microwaves.⁸⁹

Neurotransmitters

Both schizophrenia and microwave exposure involve brain dopamine alterations. Many have long attributed positive schizophrenic symptoms to dopamine increases based on differential drug effects.⁹⁰ However, negative symptom schizophrenic findings from dopamine metabolite, dopamine receptor, and drug studies indicate decreased dopamine.⁹¹ Based on behavioral changes, drug study results, and enzyme alterations, microwave exposure also indicates decreased dopamine.⁹²

Other neurotransmitter alterations correspond in both microwave bioeffect and schizophrenia. Brain postmortem tissue analysis, cerebrospinal fluid, and drug studies find decreased schizophrenic serotonin.⁹³ Although rat serotonin metabolite ratios indicate increased serotonin turnover rates on acute microwave exposure (3.1 X US pop. std.),⁹⁴ brain serotonin decrease occurs on prolonged exposure (near US occup. std.) (45, Hermann) Rat microwave exposure from birth to 15 days decreased serotonin in adults (near ½ US occup. std.)⁹⁵ Cortical synaptosome γ-aminobutyric acid (GABA) uptake and release is reported decreased in schizophrenics, who have decreased GABA neurons,⁹⁶ and synthetic enzymes. GABA receptor binding (by ³H-muscimol) decreases in rat neocortex on microwave irradiation (2.6 X US occup. std.)⁹⁷ Cholinergic system disruption impairs memory and attention; prominent schizophrenia features, however though acetylcholine alterations are indicated, evidence for either an overall increase or decrease is

inconclusive.⁹⁸ Similarly indeterminate is the microwave net effect on acetylcholine. Microwaves inactivate acetylcholine esterase activity⁹⁹ (which may increase cholinergic activity, though disputed as to the enzyme's Km), (45, Hermann) and abolish scopolamine anticholinergic effects.¹⁰⁰ However, acute rat microwave exposure decreases sodium dependent choline uptake, the rate limiting step in acetylcholine synthesis, especially in frontal cortex followed by the striatum on either pulsed or continuous wave, but only pulsation decreased hippocampal choline uptake (60 % of US pop. std.)^{101, 102} (18, Lai, 89)

The hippocampus and striatum are limbic structures-- a brain system prominent in schizophrenia pathogenesis. This system is also involved in microwave bioeffects.^{103, 104} Microwave exposure amplification of hippocampus theta rhythm is observed (30 % of US pop. std.),¹⁰⁵ with histologic and anatomic alteration reported.^{106, 107}

Hormones

Corticotrophin is indicated to mediate microwave stress,^{108, 109} and microwaves influence adrenal steroids. Satellite station operator microwave exposures produce a stress reaction of urinary increases in 11-oxycorticosteroids and stress hormone diurnal pattern shift (1/10th of US pop. std.)¹¹⁰ Rat microwave exposure yields adrenal activation resulting in adrenal medulla epinephrine and corticosteroid depletion (1.8 X US occup. std.)¹¹¹ Female rat microwave exposure increased corticosterone and ACTH, with decreased estradiol independent of pregnancy.^{112, 113, 114} Schizophrenic patients have increased cortisol with less dexamethasone cortisol suppression than controls,^{115, 116} and corticosterone increase is reported.¹¹⁷ Schizophrenics have such hypothalamic-pituitary-adrenal axis over activity with ACTH increase as to feature the metabolic syndrome.¹¹⁸ Patient cortisol lacks sleep inhibition, and correlates with paranoia and hallucination. Some negative syndrome schizophrenics have decreased melatonin.¹¹⁹ Electromagnetic fields diminish melatonin in animals.¹²⁰ (95, Frey, 94) Human melatonin decrease is both at lower frequencies,^{121, 122, 123, 124} and on cell phone use.¹²⁵ The pineal gland synthesizes melatonin from serotonin,¹²⁶ also decreased as above. Abnormal EEG and decreased melatonin are associated with pineal calcification,¹²⁷ which has lower incidence in undeveloped societies¹²⁸ who also show better schizophrenic prognosis.¹²⁹

Mitochondria Changes

Mitochondria are altered in both schizophrenia and microwave exposure. Mitochondria deformation, size reduction, and decrease in number from 20-33% in schizophrenia brain are observed.¹³⁰ Cytochrome c oxidase, of the mitochondria oxidative phosphorylation system, is reduced from 30-63% in the schizophrenic brain.¹³¹ Schizophrenic mitochondria gene expression is decreased in five pathways.¹³² Acute microwave exposure evidences mitochondria matrix density decrease, and cristae degeneration <u>in vitro</u> (1.2 X US occup. std.)¹³³ Adenosine triphosphate (ATP) and creatine phosphate (CP) levels depend on oxidative phosphorylation, which requires electron transport components of mitochondria cristae. Very brief (5 min) whole body microwave exposure significantly decreased rat brain ATP and CP levels (2.5 X occup. std.)^{134, 135}

Lipid Phosphorylation

Schizophrenic brain magnetic resonance spectroscopy shows decreased phosphomonoesters, and increased phosphodiesters. This represents reduced lipid membrane building blocks, and increased lipid degradation products. Microwave exposed rabbits decrease P³² incorporation into brain lipids (1.8 X US pop. std.)¹³⁶

Blood Brain Barrier Permeability

Molecular and cellular evidence suggests blood-brain barrier (BBB) impairment in 18-29% of Schizophrenics.¹³⁷ Non-thermal microwave alteration of the BBB permeability is consistently observed,^{138, 139, 140, 141} and is attributed to pinocytosis.^{142, 143} The alteration is proposed induced by heat shock protein phosphorylation,¹⁴⁴ and heat shock protein antibodies are among the evidence for schizophrenia BBB impairment. Studies not showing the effect have utilized short exposures, thermal microwave levels, and are criticized for procedure or publication behavior¹⁴⁵ Thermal microwave BBB studies are complicated by decreased BBB permeability at about 40° brain temperature,¹⁴⁶ but at 2° higher permeability greatly increases.^{147, 148}

Immune Alterations

A schizophrenia autoimmune etiology is indicated by several immune alterations, including abnormally high autoantibodies against brain and somatic antigens,^{149, 150} Higher autoimmune disease prevalence in these patients and their relatives is reported.^{151, 152} Foreign abstracts indicate microwaves cause autoimmune stimulation.^{153, 154, 155}

Increases of the cytokine interleukin-6 (IL-6) are a feature of autoimmune disease. (155, Ganguli) Ten reports of an increase in IL-6 in schizophrenia are versus six reporting a normal level, while four reports of an increase of IL-1_{β} in the disease are versus six reporting a normal level.¹⁵⁶ Electromagnetic field exposure of human monocytes, the most important producer these cytokines, dramatically increased production of IL-6 and IL-1_{β}.¹⁵⁷ These fields were lower in frequency than microwave.

High tumor necrosis factor (TNF) levels are reported in schizophrenia. (154,

Gaughran) Very low intensity microwave whole body exposure increases TNF production in peritoneal macrophages and spleen T cells (2 X 10⁻⁴ of US pop. std.)^{158, 159} Microwave exposure TNF increase has several other reports.^{160, 161, 162}

The balance of evidence shows B lymphocyte increase in schizophrenia (5 reports of increase versus 3 of normal levels.) (142, Rothermundt) Whole body microwave exposure increases the proportion of mouse spleen B lymphocytes.^{163, 164} This increase is not caused by proliferation, but from stimulation of already existing precursor B cell maturation,¹⁶⁵ and is under genetic control,^{166, 167} with apparent humoral mediation.¹⁶⁸ Microwaves also induce human lymphocyte lymphoblastoid transformation <u>in vitro</u>.¹⁶⁹

Anatomy and Histology

Schizophrenia reduction of medial temporal lobe structures, the hippocampalamygdala complex, is observed in 74 % of magnetic resonance imaging studies, with left lateralized findings.^{170, 171} (37, Kasai) Microwaves affect both the hippocampus and the cortex. Chinese hamster 15 day microwave exposure produces pyknotic neurons in the hippocampus, hypothalamus, and unspecified cortex areas (1.8 X US occup. std.) (110, McKee) Rat pre- thru post-natal ultra-wideband microwave exposure increased hippocampus lateral length. (111, Cobb) Such enlargement may indicate edema, reflecting pathology resulting in eventual size reduction. These rat pups stress vocalized more, and later mated less.

The thalamus is volume decreased in 42 % of schizophrenia studies, (175, Shenton) with lower neuron number in the anterioventral nucleus observed. Light and electron microscopy of hamster 22 day microwave exposure reveals cytoplasm vacuolization and chromatolysis with a pale frothy cytoplasm in ventral thalamic neurons, and little rough endoplasmic reticulum, with very few polyribosomes (3 X occup. std.)¹⁷² Dendrites had vacuoles, myelin figures, and few microtubules.

Schizophrenia cerebellum changes are evident in numerous studies of neurological signs, postmortem specimens, and in 31 % of neuroimaging studies. (175, Shenton) Atrophy is the main anatomic observation, but several studies show Purkinje cell loss. Rat and quail pre-natal prolonged microwave exposure produces Purkinje cell loss and histologic change respectively (1.2 X US occup. std. & 3.1 X US pop. std.)^{173, 174} Rat post-natal microwave exposure also produces Purkinje cell decrease and cellular changes (1.2 X US occup. std.)¹⁷⁵ (?181, Albert) Pulsed microwave rat balancing ability deficit suggests cerebellum motor influence (23 % of US pop. std.) (91, Frey, 77)

Prefrontal and parietal lobe volume reduction is reported by 60 % of studies for each area. (175, Shenton) Several microwave reports are of unspecified brain area change. Prolonged microwave rat exposure produces neuronal cytoplasm vacuolation, swelling, and beading of axons, with dendrite spine decrease (less than US occup. std.) Extended microwave exposure produces myelin degeneration in rat brain, (184, Lai) and in guinea pig or rabbit cortex (1.75 & 2.5 X US pop. std.)¹⁷⁶

A neurodevelopment schizophrenia hypothesis is favored, since autopsied brain has no inflammation or scarring. Yet, brain atrophy by apoptosis lacks gross change. Microwaves <u>in vitro</u> produce apoptosis in the Fas pathway (3.1 X US pop. std.)¹⁷⁷

Metabolic Activation

Glucose uptake and blood flow in the hallucinating brain show temporal lobe activation over baseline or control in 85 % of studies.¹⁷⁸ The temporal lobe superior gyrus is activated in some 40 % of studies, but middle gyrus or medial temporal regions of hippocampus or amygdala are often activated, with thalamus activation in some studies. (187, Weiss) Acute hearing effect pulsed microwave exposure increased rat brain glucose metabolism by [¹⁴C] 2-deoxy-D-glucose with particular prominence in the lateral geniculate, medial geniculate, the ventral medial thalamus, and in limbic structures of the mammillary bodies, and amygdala (30% of & 1.2 X US occup. std.) Only more prominent uptake by non-auditory structures was noted, with both geniculate bodies being part of the thalamus, while the mammillary bodies have a terminal hippocampal tract, and are too small for imaging. ¹⁷⁹ Rat blood flow increases significantly in the temporal cortex, lateral and medial geniculate bodies with acute microwave exposure pulsed for the hearing effect (1.6 X US occup. std.)¹⁸⁰ Therefore microwave studies particularly correspond in temporal, thalamus, and amygdala regions to those of hallucination.

Brief human cell phone¹⁸¹ and rat microwave exposure also increase brain blood flow (1.2 X US occup. std.),¹⁸² but longer exposure of pregnant rats exhibited decreased uteroplacental circulation (1.2 X US pop. std.) (117, Nakamura) (118, Yoshida) Acute psychosis studies have shown increased global brain blood flow,^{183, 184} with psychosis and delusion correlation, yet the chronic patients most studied show hypoperfusion. Microwave exposures inducing thermal effects initially increases, but eventually decreases brain blood flow, though associated with cellular injury.¹⁸⁵ Specific cerebral blood flow regions are increased while hallucinating, but sensory stimuli and endogenous verbal imagery activates brain regions of hallucinators less than non-hallucinators.^{186, 187} (187, Weiss)

Brain activation changes are widely noted in schizophrenia, particularly in the frontal lobes.¹⁸⁸ At rest, schizophrenics exhibit lower glucose utilization in the frontal lobes relative to either occipital or whole brain.¹⁸⁹ The schizophrenia prefrontal blood flow is particularly deficient while performing tasks specific to this region. (197, Taylor) Schizophrenia brain perfusion during tasks includes globally increased blood flow, or less dominant hemisphere activity and more non-dominant increases than controls. Micro-wave deficits in frontal choline uptake, maze performance, contingent negative variation, and frontal neuropsychiatric symptoms above are consistent with a prefrontal deficit.

Defects in brain area volume, mitochondria, and neurotransmitters provide basis for decreased activity in schizophrenia. Corresponding defects with microwaves and the shift of brain activity to other brain areas could have mechanism in technologic assault. Although perceptual processing is normally lateralized to the left hemisphere, pitch discrimination, non-verbal, and degenerate sounds activate the right hemisphere in health.¹⁹⁰ Microwave activation may be akin to non-verbal or degenerate sound.

A microwave mechanism for EEG delta wave increase has been proposed by corpus callosum tract fatigue, making unavailable this interhemispheric connection, with inherent corticospinal and spinocortical tract delta rhythm predominant.¹⁴⁵ Schizophrenia corpus callosum dysfunction¹⁹¹ and decreased brain area activity may enlist abnormal brain area activation. One model of gamma wave distribution relates delta wave amplitude and cortical metabolic rate in normal development to transient neuronal organizational state.¹⁹² A re-organizational state may apply in technologic assault.

Positive Symptoms

Although microwave bioeffects are consistent with negative schizophrenic symptoms,^c internal voice transmission effects provide basis for most prominent positive schizophrenic symptoms.^d Presently, casual discussion of this presentation is considered delusional by psychiatric prejudice, without detailing extensive references. Because internal voice is similar to thought, and may be directive, these technologies are capable of

^c Alogia, affective blunting, anhedonia/asociality, avolution//apathy, and attention impairment.

^d Hallucination, delusions, positive thought disorder (e.g. derailment, tangentially, incoherence, etc) bizarre behavior, and inappropriate affect.

altering thought itself and ongoing behavior. Presently, positive symptoms of attention deficit and thought disorder are explained by hallucination. Exacerbating both these symptoms would be microwave impaired working memory, and EEG entrainment capability. Microwave manipulation, then could account for the major positive schizophrenic symptoms of hallucination, delusion, attention deficit, and thought disorder.

Belief of technologic assault is most consistent with the paranoid schizophrenia subtype. More studies of this diagnosis show less genetic association, a later onset, and an increase of this form within the past century is reported.^{193, 194} Schizophrenia is apparently preponderantly sporadic,^{195, 196} with EEG abnormalities reported as more frequent, in this form.¹⁹⁷ Although first admission studies have indicated a decline in schizophrenia, changing demographic and diagnostic patterns question true incidence change,¹⁹⁸ with diagnoses of borderline states,¹⁹⁹ and paranoid psychosis²⁰⁰ matching some declines, while a recent review shows a schizophrenia incidence increase.²⁰¹

Ocular Disease

Microwave exposures are known to produce eye disease. Subcapsular cataracts are particularly produced by microwaves.^{202, 203} Anterior subcapsular cataracts were significantly more prevalent in schizophrenics than a visually impaired population, without medication association, except that phenothiazines actually had less cataract prevalence.²⁰⁴ As expected for a group of little occupational exposure, schizophrenics have less cataract incidence, of all types, than the general population,²⁰⁵ but schizophrenia cataracts have been associated with high doses of chlorpromazine (a phenothiazine.)²⁰⁶ Microwave exposures have occupationally been associated with retinopathy,²⁰⁷ (30, Hanson) and experimentally produce retinal damage.²⁰⁸ Schizophrenia retinopathy is associated with thioridazine, ²⁰⁹ and

generally with phenothiazines.²¹⁰ (217, Bond) All the schizophrenia ocular disease associated drugs are older, generically available, and may have public medical assistance or patient profile prescriptive preferences. Phenothiazines were so broadly utilized that direct association with schizophrenia cannot be excluded.

Standards and Environmental Considerations

East European and Russian occupational microwave standards of 10 μ W/cm² are based on a neurasthenia syndrome.²¹¹ Reported symptoms are headache, dizziness, increased irritability, loss of appetite, sleepiness, increased fatigability, sweating, difficulties in concentration or memory, depression, emotional instability, dermatographism, thyroid enlargement, and tremor of the extended fingers. (49, Silverman) The American study of increased human EEG delta waves noted short-term memory impairment, concentration inhibition, irratibility, apprehension, frontal headache, and such sluggishness as to interfere with work the next day. (48, Bise) This syndrome is consistent with many schizophrenia symptoms.

The Russian standard has contrasted with a 1000 times greater US standard of 10 mW/cm², which was too weakly written to sustain lawsuit. (222, Steneck) The original US standard was set at one-tenth the level known to increase body temperature. The main microwave research sponsor, the Defense Department has vigorously defended this thermal rationale with suppression of non-thermodynamic effect investigations.²¹² Standard setting for optimal equipment performance on national security grounds is suggested.²¹³ Present US standards (ANSI/IEEE C95.1) lowered the occupational standards within certain frequencies, and finally set population standards, though at ~100 times the Russian. (14,

Ghandi) There are many reported effects at, or near these standards, which are certainly not so strict as to exclude all effects, however detrimental.

A 1975 Environmental Protection Agency survey indicated that less than 1% of the population was routinely exposed to more than 1 μ W/cm², and that high exposure areas (building tops with radiofrequency transmitter clusters) could run as high as 100-200 μ W/cm². (222, Stencek) Cell phones can reach 200 mW power output.²¹⁴ Although few microwave correlates of schizophrenia are at, or below these levels, neither well studied is chronic exposure, with considerable exposurechange since 1975.

Unknown is an environmental microwave relationship to schizophrenia, except for those correlations here reviewed. Even though a manufactured system may meet the standards, sources are proliferating, and standards may be exceeded in many situations, particularly with increasing cell phone use. Potentially toxic effects are in cell phone reports, and base station proximity increases risk, with recognized over exposure by heat-sealing appliances. (222, Steneck) Dermatologic electromagnetic hypersensitivity syndromes are reported by patients, as well as a type resembling neurasthenia recognized by the Russians.²¹⁵ Though such syndromes are unconfirmed, yeast cell effects are some seven orders of magnitude below the Russian standard.²¹⁶ Microwaves are a proposed mechanism for a reported sunspot activity association with schizophrenia.²¹⁷

A schizophrenia neurodevelopment hypothesis is now favored, but there is evidence for a neurodegenerative process in a sub-population.^{218, 219} Neurodegenerative diseases such as amyotrophic lateral sclerosis (ALS), Alzheimer's, and Parkinsonism are linked to electromagnetic field exposure.²²⁰ For ALS, data indicating increased risk is relatively strong, while for Alzheimer's the data is weaker than for ALS, and Parkinsonism is only linked with little evaluative data.

SUMMARY AND CONCLUSION

Microwave technology is capable of internal voice transmission. Microwave internal voice weapons are considered²²¹ (7, DIA) (3, Oskar) and a weapon has been referenced, (9, Army) with the basic technology described. (2, Justensen) (5, Burnkan) Continuous symptoms present in psychosis can be maintained by available tracking technology. Since similar means are a frequent patient complaint, it is compulsory that methods be developed to rule out involvement of these technologies in delusional disorder and psychosis. It is unethical to further ignore the evidence and disrespect these patients.

Microwave bioeffects have a high level of congruence with major lines of schizophrenia investigation. In both schizophrenia and microwave exposure, there is cognitive deficit, and a number of electrophysiologic signs including decreased contingent negative variation, decreased auditory event related response, and increased EEG delta waves. Startle response and galvanic skin response is found decreased in both conditions. In the neurotransmitter levels of both conditions, serotonin is found decreased, with dopamine and GABA indicated as decreased. The limbic system is afflicted in both situations. Hormone changes of melatonin decrease, and adrenal activation are common to both conditions. Immune function, mitochondria, and the blood-brain barrier are reported similarly altered in both situations. Microwaves induce deleterious histology in several brain structures observed reduced in schizophrenia. Microwave exposure activates brain structures corresponding to those noted on hallucination. Subcapsular cataracts have been associated with both conditions. Retinopathy is associated with both widely prescribed anti-psychotic medication, and microwave exposure. Microwave voice transmission, and EEG entrainment provide basis for positive symptoms. The almost comprehensive congruence between microwave bioeffects and schizophrenia may not apply to all patients, but is most consistent with the negative symptom group.

The congruence of microwave bioeffects with schizophrenia symptoms does not have to involve voice transmittal in a technologic etiology. Potentially toxic effects to functioning exist near, and at exposure standards. Hypersensitivity syndromes are reported at lower frequencies than microwave, although any syndrome has been unconfirmed. Neurodegenerative diseases are also associated with lower frequency exposure, particularly ALS and Alzheimer's. The potential for voice transmittal to mimic positive schizophrenia symptoms, and the congruence of other symptoms with microwave bioeffects indicates that a technologic etiology may involve more than a few patients.

The medical community has been remiss in refusing investigation of such an etiology. Psychiatrists have actively ignored longstanding patient complaints of being affected as basically herein referenced. Microwave bioeffects, including sound and voice perception have long been described. The evidence for a technologic etiology regarding microwaves practically comprehensively correlates with schizophrenic symptoms to such congruence that a mathematic congruent state cannot be excluded. This hypothesis is more defined than any other means of pathogenesis, and should mandate investigation to develop methods for ruling out such an etiology. Though direct proof is lacking as to specific cases, investigation must begin. Of course the hypothesis may not involve all cases, for multifactor etiologies are common to medicine, and reference is often made to "the schizophrenias."

Patients subject to internal voice assault would have hallucination, and likely paranoia with belief that voices are transmitted to them. It would be most probable among

sporadic cases with non-adolescent onset, having some or all of the correlations here noted. Clinical investigation would include radiofrequency measurement. Attention should be given to likely cranial directional localization within the specified Brunkan patent hearing spectrum. Establishing radiation characteristics with the Brunkan patent burst and pulse pattern or multiple frequencies as in the Stocklin patent would also be highly pertinent, but less important.

Investigation of responses within and outside of rooms shielded from electromagnetic radiation is relevant. Practical considerations are that shielded facilities already exist for MRI and magnetoencephalgraphy. Observations of hallucination, event related auditory response, contingent negative variation, or EEG delta wave index in selected patients would likely be parameters more immediately responsive to microwave cessation. Although existing facilities may be adequately shielded,²²² the shielding must be radar effective, with serious determination of adequacy.

Subcapsular cataract and retinopathy epidemiologic study in schizophrenia would also have relevance. The specific cataract type is known to be microwave induced, and is reported without medication association. Patient signs relating to other microwave bioeffects would have bearing on any coincidence of these symptoms, which was indeterminate to this review, and is pertinent to a technologic etiology.

Acknowledgements: Thanks are given to God for inspiration, and to Dr. Paul Canner for suggestions.

Patents are printable free from the U. S. Patent Office website.

REFERENCES

¹ Frey AH, "Human Auditory System response to modulated electromagnetic energy" J Applied Physiol 17(4): 689-92, 1962. <u>http://www.raven1.net/frey.htm</u>

Justesen DR. "Microwaves and Behavior" Am Psychologist 392(Mar): 391-401, 1975. Excerpted reference at <u>http://www.raven1.net/v2succes.htm</u>

² Oskar KJ. "Effects of low power microwaves on the local cerebral blood flow of conscious rats" Army Mobility Equipment Command Report, NASA Technical Report. # AD-A090426, 1980. Abstract at http://www.raven1.net/v2s-nasa.htm

³ Kohn B. "Communicating Via the Microwave Auditory Effect" Defense Department Awarded SBIR Contract # F41624-95-C9007, 1993. at <u>http://es.epa.gov/ncer_abstracts/sbir/other/monana/kohn.html</u> & <u>http://www.raven1.net/v2s-kohn.htm</u>

Brunkan WB. Patent #4877027 "Hearing system" USPTO granted 10/31/89.

⁴ Stocklin PL. Patent #4858612 "Hearing device" USPTO granted 8/22/89.

"Surveillance Technology, 1976: policy and implications, an analysis and compendium of materials: a staff report of the Subcommittee on Constitutional Rights of the Committee of the Judiciary. United States Senate, Ninety-fourth Congress, second session p 1280, US GOV DOC Y 4.J 882:SU 7/6/976.

⁵ Castelli CJ. "Questions Linger about Health Effects of DOD's 'Non-Lethal Ray' Inside the Navy 14(12): 1-6, 2001. http://globalsecurity.org/org/news/2001/e20010327questions.htm

http://www.pjproject.org/usaf.html

⁶ Center for Army Lessons Learned Thesaurus at <u>http://call.army.mil/products//thesaur/00016275.htm</u>

⁷ McMurtrey J. Technologies of 'Hallucination' and Behavioral Influence'' Companion paper.

⁸ Flaum M and Schultz SK. "The Core Symptoms of Schizophrenia" Ann Med 28(6): 525-31, 1996.

⁹ Isselbacher KJ, Adams RD, Brunwald E, Petersdorf RG (eds.) and Wilson JD. <u>Harrison's Principles of Internal</u> <u>Medicine</u> Ninth Ed., McGraw-Hill, New York, p. 150, 1980.

¹⁰ American Psychiatric Association DSM-IV Task Force. <u>Diagnostic and Statistical Manual of Mental Disorders Fourth</u> <u>Edition (DSM-IV-TR[™])</u> American Psychiatric Association, p 297-343, 2000.

¹¹ Gandhi OP. "Electromagnetic Fields: Human Safety Issues" Ann Rev Biomed Eng 4: 211-34, 2002.

¹² Heinrichs RW and Zakzanis KK. "Neurocognitive Deficit in Schizophrenia: A Quantitative Review of Evidence" Neuropsych 12(3): 426-45, 1998.

¹³ Stip E. "Memory Impairment in Schizophrenia: Perspectives form Psychopathology and Pharmacotherapy" Can J Psychiatry 41(8 Suppl. 2): S27-S34, 1996.

¹⁴ Goldman-Rakic PS. "Working Memory Dysfunction in Schizophrenia" J Neuropsychatr 6(4): 348-55, 1994.

¹⁵ Lai H, Carino MA, Horita A, and Guy AW. "Low-Level Microwave Irradiation and Central Cholinergic Activity: A Dose Response Study" Bioelectromagnetics 10: 203-8, 1989.

¹⁶ Lai H, Horita A, and Guy AW. "Microwave Irradiation Affects Radial-Arm Maze Performance in the Rat" Bioelectromagnetics 15: 95-104, 1994.

¹⁷ Wang B and Lai H. "Acute Exopsure to Pulsed 2450-MHz Microwaves Affects Water-Maze Performance of Rats" Bioelectromagnetics 21: 52-6, 2000.

¹⁸ D'Andrea JA, DeWitt JR, Gandhi OP, Stensaas S, Lords JL, and Nielson HC. "Behavioral and Physiological Effects of Chronic 2,450 MHz Microwave Irradiation of the Rat at 0.5 mW/cm²" Bioelectromagnetics 7: 45-56, 1986.

¹⁹ D'Andrea JA, DeWitt JR, Emmerson RY, Bailey C, Stensaas S, and Gandhi OP. "Intermittent Exposure of Rats to 2450 MHz Microwaves at 2.5 mW/cm²: Behavioral and Physiological Effects" Bioelectromagnetics 7: 315-28, 1986.

²⁰ McRee DI, Elder JA, Gage MI, Reiter LW, Rosenstein LS, Shore ML, Galloway WD, Adey WR, and Guy AW. "Effects of Nonioniziong Radiation on the Central Nervous System, Behavior, and Blood: A Progress Report" Environ Health Perspect 30: 123-31, 1979.

²¹ Thomas JR, Finch ED, Fulk DW, and Burch LS. "Effects of Low-Level Microwave Radiation on Behavioral Baselines" Ann N Y Acad Sci 247: 425-32, 1975.

²² Gage MI. "Behavior in Rats After Exposures to Various Power Densities of 2450 MHz Microwaves" Neurobehavioral Toxicol 1: 137-43, 1979.

²³ Thomas JR, Schrot J, and Banvard RA. "Comparative Effects of Pulsed and Continuous-Wave 2.8-GHz Microwaves on Temporally Defined Behavior" Bioelectromagnetics 3: 227-35, 1982.

²⁴ Schrot J, Thomas JR, and Banvard RA. "Modification of the Repeated Acquisition of Response Sequences in Rats by Low-Level Microwave Exposure" Bioelectromagnetics 1: 89-99, 1980.

²⁵ Thomas JR, Veandle SS, and Burch LS. "Modification of Internal Discriminative Simulus Control of Behavior by Low Levels of Pulsed Microwave Radiation" In: Johnson CC and Shore ML (eds.) <u>Biological Effects of Electromagnetic Waves</u> HEW Publications (FDA) 77-8010, Rockville, MD, p 201-14, 1976.

²⁶ Isa AR. "Non-Ionizing radiation exposure causing ill-health and alopecia areata" Med J Malaysia 40(3): 235-8, 1991.

²⁷ Hansson HA. "Effects on the Nervous System by Exposure to Electromagnetic Fields: Experimental and Clinical Studies" In: <u>Electromagnetic Fields and Neurobehavioral Function</u> Prog in Clin and Biol Res 257: 119-34, 1988.

²⁸ Klein C, Heinks T, Andresen B, Berg P, and Moritz S. "Impaired Modulation of the Saccadic Contingent Negative Variation Preceeding Antisaccades in Schizophrenia" Biol Psychiatry 47: 978-90, 2000.

²⁹ van den Bosch. "Contingent Negative Variation and Psychopathology: Frontal-Central Distribution, and Association with Performance Measures" Biol Psychiatry 18(6): 615-34, 1983.

³⁰ Abraham P, McCallum WC, and Gourlay J. "The CNV and Its Relation to Specific Psychiatric Syndromes" In: McCallum WC and Knott JR (eds.) <u>The Responsive Brain</u> Wright, Bristol UK, p 144-49, 1976.

³¹ Timsit-Berthier M, Gerono A, Rousseau JC, Mantanus H, Abraham P, Verhey EHM, Lamers T, and Emonds P. "An International Pilot Study of CNV in Mental Illness, Second Report" In: Karrer R, Cohen J, and Tuering P (eds.) <u>Brain and Information</u> New York: New York Academy of Science, p 629-37, 1984.

³² Freude G, Ullsperger P, Eggert S, and Ruppe I. "Effects of Microwaves Emitted by Cellular Phones on Human Slow Brain Potentials" Bioelectromagnetics 19: 384-7, 1998.

³³ Freude G, Ullsperger P, Eggert S, and Ruppe I. "Microwaves emitted by cellular telephones affect human slow brain potentials" Eur J Appl Physiol 81: 18-27, 2000.

³⁴ Kasai K, Iwanami A, Yamasue H, Kuroki N, Nakagome K, and Fukuda M. "Neuroanatomy and neurophysiology in schizophrenia" Neurosci Res 43: 93-110, 2002.

³⁵ O'Donnell BF, McCarley RW, Potts GF, Salisbury DF, Nestor PG, Hiryasu Y, Niznikiewicz MA, Barnard J, Shen ZJ, Weinstein DM, Bookstein FL, and Shenton ME. "Identification of neural circuits underlying P300 abnormalities in schizophrenia" Psychophysiology 36: 388-98, 1999.

³⁶ McCarley RW, Faux SF, Shenton ME, Nestor PG, and Adams J. "Event-related potentials in schizophrenia: their biological and clinical correlates and a new model of schizophrenic pathophysiology" Schizophrenia Res 4: 209-31, 1991.

³⁷ Tiihonen J, Hari R, Naukkarinen H, Rimon R, Jousmaki V, and Kajola M. "Modified Activity of the Human Auditory Cortex During Auditory Hallucinations" Am J Psychiatry 149(2): 255-7, 1992.

³⁸ Hari R and Makela JP. "Modification of neuromagnetic responses of the human auditory cortex by masking sounds" Exp Brain Res 71: 87-92, 1988.

³⁹ Sagalovich BM and Melkumova GG. "[Research on the action of superhigh-frequency electromagnetic waves on evoked potentials of auditory centers in connection with prospects for using inadequate auditory stimulation]" Vestnick Otorinolaring. 4: 3-8, 1974. (An English translation is available in Popov, SL (ed.) "Effects of Non-Ionizing Electromagnetic Radiation" JPRS report # 64532, Arlington, VA, p. 23-30, 1975.)

⁴⁰ Guy AW, Chou CK, Lin JC, and Christensen D. "Microwave-Induced Acoustic Effects in Mammalian Auditory Systems and Physical Materials" Ann N Y Acad Sci 247: 194-217, 1975.

⁴¹ Seaman RL and Lebovitz RM. "Thresholds of Cat Cochlear Nucleus Neurons to Microwave Pulses" Bioelectromagnetics 10: 147-60, 1989.

⁴² Hermann DM and Hossman K-A. "Neurological effects of microwave exposure related to mobile communication" J Neurol Sci 152: 1-14, 1997.

⁴³ Michaelson SM. "Sensation and Perception of Microwave Energy" In: Michaelson SM, Miller MW, and Carstensen EL (eds.) <u>Fundamental and Applied Aspects of Nonionizing Radiation</u> Plenum Press, New York, p 213-29, 1975.

⁴⁴ Ford JM. "Schizophrenia: the broken P300 and beyond" Psychophysiology 36: 667-82, 1999.

⁴⁵ Bise W. "Low Power radio-frequency and microwave effects on human electroencephalogram and behavior" Physiol Chem Phys 10(5): 387-98, 1978.

⁴⁶ Silverman C. "Nervous And Behavioral Effects of Microwave Radiation in Humans" J Epidemol 97: 219-24, 1973.
⁴⁷ Reiser H-P, Dimpfel W, and Schober F. "The Influence of Electromagnetic Fields on Human Brain Activity" Eur J Med Res 1: 27-32, 1995.

⁴⁸ Baranski S and Edelwejn Z. "Electroencephalographic and Morphological Investigations of the Influence of Microwaves on the Central Nervous System" Acta Physiol Pol 18(4): 423-36, 1967.

⁴⁹ Shandala, MG, Dumanskii UD, Rudnev MI, Ershova LK, and Los IP. "Study of Nonionizing Microwave Radiation Effects upon the Central Nervous System and Behavior Reactions" Environ Health Perspect 30: 115-21, 1979.

⁵⁰ Voroh'ev VV, Konovalov VF, Gorelkova TF, and Gal'chenko AA. "[The Electrical activity of symmetrical areas of the rat cerebral cortex during the use of a low-intensity UHF field]" Fiziol Zh Im I M Sechenova 80(12): 55-61, 1994.

⁵¹ Takashima S, Onaral B, and Schwan HP. "Effects of Modulated RF Energy on the EEG of Mammalian Brains: Effects of Acute and Chronic Irradiation" Rad and Environ Biophys 16: 15-27, 1979.

⁵² Persinger MA, Richards PM, and Koren SA. "Differential Entrainment of Electroencephalographic Acitvity by Weak Complex Electromagnetic Fields" Percept Motor Skills 84: 527-36, 1997.

⁵³ Karson CN, Coppola R, Morihisa JM, and Weinberg DR. "Computed Electroencephalographic Acitvity Mapping in Schizophrenia: The Resting State Reconsidered" Arch Gen Psychiatry 44: 514-17, 1987.

⁵⁴ Miyauchi T, Tanaka K, Hagimoto H, Miura T, Kishimoto H, and Matsushita M. "Computerized EEG in Schizophrenic Patients" Biol Psychiatry 28: 488-94, 1990.

⁵⁵ Morihisa JM, Duffy FH, and Wyatt RJ. "Brain Electrical Activity Mapping (BEAM) in Schizophrenic Patients" Arch Gen Psychiatry 40: 719-28, 1983.

⁵⁶ Nagase Y, Okubo Y, Marsuura M, Kojima T, and Toru M. "EEG Coherence in Unmedicaated Schizophrenic Patients: Topographical Study of Predominantly Never Medicated Cases" Biol Psychiatry 32: 1028-34, 1992.

⁵⁷ Takizawa Y, Wada Y, Horita M, Kitazawa S, Futamata H, and Hashimoto T. "Quantitative analysis of EEG background activity in drug naïve schizophrenic patients" Rinsho Byori 42(7): 759-63, 1994.

⁵⁸ Miyauchi T, Endo S, Kajiwara S, Ishii M, and Okajima J. "Computerized electroencephalogram in untreated schizophrenics: A comparison between disorganized and paranoid types" Psychiatr Clin Neurosci 50: 71-8, 1996.

⁵⁹ Begic D, Hotujac L, and Jokic-Begic N. "Quantitative EEG in 'positive' and 'negative' schizophrenia" Acta Psychiatr Scand 101: 307-11, 2000.

⁶⁰ Gladerisi S, Mucci A, Mignone ML, Maj M, and Kemali D. "CEEG mapping in drug-free schizophrenics: Differences from healthy subjects and changes induced by haloperidol treatment" Schizophr Res 6: 15-24, 1992.

⁶¹ Pascaul-Marquai RD, Lehmann D, Koenig T, Kochi K, Merlo MCG, Hell D, and Koukkou M. "Low resolution brain electromagnetic tomography (LORETA) functional imaging in acute, neuroleptic-naive, first episode, productive schizophrenia" Psychiatr Res Neuroimaging Sect 90: 169-79, 1999.

⁶² Fenton GW, Fenwich PRC, Dollimore J, Dunn TL, and Hirsch SR. "EEG Spectral Analysis in Schizophrenia" Brit J Psychiatr 136: 445-55, 1980.

⁶³ Guenther W, Breitling D, Banquet JP, Marcie P, and Rondot P. "EEG Mapping of Left Hemisphere Dysfunction during Motor Performance in Schizophrenia" Biol Psychiatry 21: 249-62, 1986.

⁶⁴ Guenther W, Davous P, Godet JL, Guillibert E, Breitling D, and Rondot P. "Bilateral Brain Dysfunction During Motor Activation in Type II Schizophrenia Measured by EEG Mapping" Biol Psychiatry 23: 395-411, 1988.

⁶⁵ Guenther W and Breitling D. "Predominant Sensorimotor Area Left Hemisphere Dysfunction in Schizophrenia Measured by Brain Electrical Activity Mapping" Biol Psychiatry 20: 515-32, 1985.

⁶⁶ Sponheim SR, Clementz BA, Iacono WG, and Beiser M. "Resting EEG in first-episode and chronic schizophrenia" Psychophysiol 31: 37-43, 1994.

⁶⁷ Locatelli M, De Angeli A, Leone F, Grassi B, and Scarone S. "Factor Analysis and Computerized EEG: Preliminary Data on Schizophrenic Patients" Int J Neurosci 72: 265-70, 1993.

⁶⁸ Takeuchi K, Takigawa M, Fukuzako H, Hokazono Y, Hirakawa K, Fukuzako T, Ueyama K, Fujimoto T, and Matsumoto K. "Correlation of Third Ventriclar Enlargement and EEG Slow Wave Activity in Schizophrenic Patients" Psychiatr Research: Neuroimaging 55: 1-11, 1994.

⁶⁹ Clementz BA, Sponheim SR, Iacono WG, and Beiser M. "Resting EEG in first-episode schizophrenia patients, bipolar psychosis patients, and their first-degree relatives" Psychophysiol 31: 486-94, 1994.

⁷⁰ Sengoku A and Takagi S. "Electroencephalographic findings in functional psychoses: State or trait indicators?" Psychiatr Clin Neurosci 52(4): 375-88, 1998.

⁷¹ Stevens JR, Bigelow L, Denney D, Lipkin J, Livermore AH, Rauscher F, and Wyatt RJ. "Telemetered EEG-EOG During Psychotic Behaviors of Schizophrenia" Arch Gen Psychiatry 36: 251-62, 1979.

⁷² Stevens JR and Livermore A. "Telemetered EEG in schizophrenia: spectral analysis during abnormal behavior episodes" J Neurol Neurosurg & Psychiatr 45: 385-95, 1982.

⁷³ Whitton JL, Moldofsky H, and Lue F. "EEG Frequency Patterns Associated with Hallucinations in Schizophrenia and "Creativity" in Normals" Biol Psychiatry 13(1): 123-33, 1978.

⁷⁴ Bawin SM, Gavalas-Medici RJ, and Adey WR. "Effects of Modulated Very High Frequencey Fields on Specific Brain Rhythms in Cats" Brain Res 58: 365-84, 1973.

⁷⁵ Servantie B, Servantie AM, and Etienne J. "Synchronization of Cortical Neurons by Pulsed Microwave Field as Evidenced by Spectral Analysis of Electrocorticograms from the White Rat" Ann N Y Acad Sci 247: 82-6, 1975.

⁷⁶ Bell G, Marino A, Chesson A, and Struve F. "Electrical states in the rabbit brain can be altered by light and electromagnetic fields" Brain Res 570: 307-15, 1992.

⁷⁷ Gavalas RJ, Walter DO, Hamer J, and Adey WR. "Effect of Low-level, Low-frequency Electric Fields on EEG and Behavior in Macaca Nemestrina" Brain Res 18: 491-501, 1970.

⁷⁸ Bell GB, Marino AA, and Chesson. "Frequency specific responses in the human brain caused by electromagnetic fields" J Neurol Sci 123: 26-32, 1994.

⁷⁹ Doswald-Beck L and Cauderay GC. "The development of New Antipersonnel Weapons" Int Rev Red Cross 279: Nov 1 1990. also at <u>http://www.dcn.davis.ca.us/~welsh/factsht.htm</u>

⁸⁰ Bernstein AS. "Orienting Response Research in Schizophrenia: Where We Have Come and Where We Might Go" Schizophrenia Bull 13(4): 623-41, 1987.

⁸¹ Gruzelier J. "Bilateral electrodermal activity and cerebral mechanisms in syndromes of schizophrenia and the schizotypal personality" Int J Psychophysiology 16: 1-16, 1994.

⁸² Thompson WD and Bourgeois AE. "Nonionizing Radiations" In: Furchtgott E (ed.) <u>Pharmacological and</u> <u>Biophysical Agents and Behavior</u> Academic Press, New York, London, 1971.

⁸³ Docherty NM. "Affective Reactivity of Symptoms as a Process Discriminator in Schizophrenia" J Nerv Mental Dis 184(9): 535-41, 1996.

⁸⁴ Mitchell CL, McRee DI, Peterson NJ, and Tilson HA. "Some Behavioral Effects of Short-Term Exposure of Rats to 2.45 GHz Microwave Radiation" Bioelectromagnetics 9: 259-68, 1988.

⁸⁵ Seaman RL and Beblo DA. "Modification of Acoustic Startle by Microwave Pulses in the Rat" Bioelectromagnetics 13: 323-28, 1992.

⁸⁶ Seaman RL, Beblo DA, and Raslear TG. "Modification of Acoustic and Tactile Startle by Single Microwave Pulses" Physiol Behav 55(3): 587-95, 1994.

⁸⁷ Galvin MJ, Tilson HA, Mitchell CL, Peterson J, and McRee DI. "Influence of Pre- and Postnatal Exposure of Rats to 2.45-GHz Microwave Radiation on Neurobehavioral Function" Bioelectromagnetics 7: 57-71, 1986.

⁸⁸ Frey AH. "Behavioral Effects of Electromagnetic Energy" In: Hazzard DG (ed.) <u>Symposium on Biological Effects</u> and <u>Measurement of Radiofrequency/Microwaves</u> HEW Publications (FDA), 77-8026 Rockville, MD, p 11-22, 1977.

⁸⁹ Frey AH, Feld SR, and Frey B. "Neural Function and Behavior: Defining the Relationship" Ann N Y Acad Sci 247: 433-8, 1975.

⁹⁰ Berger PA. "Biochemistry and the Schizophrenias: Old Concepts and New Hypotheses" J Nerv Mental Dis 169(2): 90-9, 1981.

⁹¹ Rao ML and Moller H-J. "Biochemical Findings of Negative Symptoms in Schizophrenia and Their Putative Relevance to Pharmacologic Treatment" Neuropsychobiology 30: 160-72, 1994.

⁹² Frey AH. "An Integration of the Data on Mechanisms with Particular Reference to Cancer" In: Frey AH (ed.) <u>On the</u> <u>Nature of Electromagnetic Field Interactions with Biological Systems</u> RG Lanes Co., Austin TX, p 9-28, 1994.

⁹³ Abi-Dargham A, Laruelle M, Aghajanian GK, Charney D, and Krystal J. "The Role of Serotonin in the Pathophysiology and Treatment of Schizophrenia" J Neuropsychiatr 9(1): 1-17, 1997.

⁹⁴ Inaba R, Shishido K, Okada A, and Moroji T. "Effects of whole body microwave exposure on the rat brain contents of biogenic amines" Eur J Appl Physiol 65: 124-8, 1992.

⁹⁵ Guessar A, Lescoat G, and Maniey A. "Influence of Postnatal Exposition to Microwaves on Brain and Hypothalmo-Pituitary Monoamines in the Adult Male Rat" Physiologie 20(2): 71-4, 1983.

⁹⁶ Lewis DA, Pierri JN, Volk DW, Melchitzky DS, and Woo T-UW. "Altered GABA Neurotransmission and Prefrontal Cortical Dysfunction in Schizophrenia" Biol Psychiatry 46: 616-26, 1999.

Benes FM and Berretta S. "GABAergic Interneurons: Implications for Understanding Schizophrenia and Bipolar Disorder" Neuropsychopharmacology 25(1): 1-27, 2001.

⁹⁷ Kolomytkin O, Yurinska M, Zharikov S, Kuznetsov V, and Zharikov A. "Response of Brain Receptor Systems to Microwave Energy Exposure" In Frey AH (ed.) <u>On the Nature of Electromagnetic Field Interaction with Biological</u> <u>Systems</u> R G Lanes Co, Austin, TX, p 195-206, 1994.

⁹⁸ Hyde TM and Crook JM. "Cholinergic systems and schizophrenia: primary pathology or epiphenomenona?" J Chem Neuroanat 22: 53-63, 2001.

⁹⁹ Kolomytkin OV, Kuznetxov VI, and Akoev IG. "Microwaves Afftect the Function of Reconstituted and Native Receptor Membranes of the Brain" In: Alen MJ, Cleary SF, and Hawkridge FM (eds.) <u>Charge and Field Effects in Biosystems –2</u> Plenum, New York & London, p 149-60, 1989.

¹⁰⁰ Monahan JC. "Microwave – Drug Interaction in the Cholinergic Nervous System of the Mouse" In <u>Progress in</u> <u>Clinical and Biological Research</u> 257: 309-26, 1988.

¹⁰¹ Lai H, Horita A, and Guy AW. "Acute Low-Level Microwave Exposure and Central Cholinergic Activity: Studies on Irradiation Paramenters" Bioelectromagnetics 9: 355-62, 1988.

¹⁰² Lai H, Horita A, Chou CK, and Guy AW. "Effects of Low-Level Microwave Irradiation on Hippocampal and Frontal Cortical Choline Uptake are Classically Conditionable" Pharm Biochem Behav 27: 635-9, 1987.

¹⁰³ Stocklin PL and Stocklin BF. "Low Power Microwave Effects on the Human Electroencephalogram: Supporting Results of Bise" Physiol Chem 13: 175-7, 1981.

¹⁰⁴ Shtemberg AS, Uzbekov MG, Shikhov SN, Bazyan AS, and Chernyakov GM. "Some Neurotropic Effects of Low-Intensity Electromagnetic Waves in Rats with Different Typological Characteristics of Higher Nervous Activity" Neurosci Behav Physiol 31(5): 547-53, 2001.

¹⁰⁵ Grigor'ev IuG, Luk'ianov SN, Makarov VP, and Rynskov VV. "[Total bioelectric activity of various structures of the brain in low-intensity microwave irradiation]" Radiats Biol Radioecol 35(1): 57-65, 1995.

¹⁰⁶ McKee A, Dorsey CH, Eisenbrandt DL, and Woden NE. "Ultrastructural Observations of Microwave-Induced Morphological Changes in the Central Nervous System of Hamsters" Bioelectromagnetics 1: 206, 1980.

¹⁰⁷ Cobb RL, Jauchem JR, Mason PA, Dooley MP, Miller SA, Ziriax JM, and Murphy MR. "Neural and Behavioral Tertaological Evaluation of Rats Exposed to Ultra-Wideband Electromagnetic Fields" Bioelectromagnetics 21: 324-37, 2000.

¹⁰⁸ Lai H, Horita A, Chou C-K, and Guy AW. "Low-Level Microwave Irradiations Affect Central Cholinergic Activity in the Rat" J Neurochem 48(1): 40-5, 1987.

¹⁰⁹ Lai H, Carino MA, Horita A, and Guy AW. "Corticotrophin-Releasing Factor Antagonist Blocks Microwave-Induced Decreases in High –Affinity Choline Uptake in the Rat Brain" Brain Res Bull 25: 609-12, 1990.

¹¹⁰ Vangelova K, Israel M, and Mihaylov S. "The Effect of Low Level Radiofrequencey Electromagnetic Radiation on the Excretion Rates of Stress Hormones in Operators During 24-hour Shifts" Cent Eur J Publ Health 10(1-2): 24-8, 2002.

¹¹¹ Parker LN. "Thyroid suppression and adrenomedulary activation by low-intensity microwave radiation" Am J Physiol 224(6): 1388-90, 1973.

¹¹² Nakamura H, Seto T, Nagase H, Yoshida M, Dan S, and Ogino K. "Effects of exposure to microwaves on cellular immunity and placental steroids in pregnant rats" Occup Environ Med 54: 676-80, 1997.

¹¹³ Nakamura H, Nagase H, Ogino K, Hatta K, and Matsuzaki I. "Uteroplacental circulatory disturbance by prostaglandin $F_{2\alpha}$ in rats exposed to microwaves" Reprod Toxicol 14: 235-40, 2000.

¹¹⁴ Yoshida Y, Seto T, Ohsu W, Hayashi S, Okarawa T, Nagase H, Yoshida M, and Nakamura H. "[Endocrine mechanism of placental circulatory disturbances induced by microwave in pregnant rats]" Nippon Sanka Fujinka Gakkai Zasshi 47(2): 101-8, 1995.

¹¹⁵ Altamura AC, Boin F, and Maes M. "HPA axis and cytokines dysregulation in schizophrenia: potential implications for the antipsychotic treatment" Eur Neuropsychopharmacology 10: 1-4, 1999.

¹¹⁶ Gispen-de Wied CC. "Stress in schizophrenia: an integrative view" Eur J Pharmacol 405: 375-84, 2000.

¹¹⁷ Conroy RTWL, Hughes BD, and Mills JN. "Circadian Rhythm of Plasma 11-Hydroxycoticosteroids in Psychiatric Disorders" Br Med J Aug 17; 2(615): 405-7, 1968.

¹¹⁸ Ryan MCM and Thakore JH. "Physical consequences of schizophrenia and its treatment: The Metabolic Syndrome" Life Sci 71: 239-57, 2002.

¹¹⁹ Sandyk R and Kay SR. "Pineal Melatonin in Schizophrenia: A Review and Hypothesis" Schizophr Bull 16(4): 653-61, 1990.

¹²⁰ Reiter RJ. "Alterations of the Circadian Melatonin Rhythm by the Electromagnetic Spectrum: A Study in Environmental Toxicology" Reg Tox Pharm 15: 226-44, 1992.

¹²¹ Pfluger DH and Minder CE. "Effects of exposure to 16.7 Hz magnetic fields on Urinary 6-hydroxymelatonin sulfate excretion of Swiss railway workers" J Pineal Res 21: 91-100, 1996.

¹²² Juutilainen J, Stevens RG, Anderson LE, Hansen NH, Kilpelainen M, Kumlin T, Laitinen JT, Sobel E, and Wilson BW. "Nocturnal 6-hydroxymelatonin sulfate excretion in female workers exposed to magnetic fields" J Pineal Res 28: 97-104, 2000.

¹²³ Burch JB, Reif JS, Yost MG, Keefe TJ, and Pitrat CA. "Nocturnal excretion of a urinary melatonin metabolite among electric utility workers" Scand J Work Environ Health 24(3): 183-9, 1998.

¹²⁴ Karasek M, Woldanska-Okonska M, Czernicki J, Zylinska K, and Swietoslawski J. "Chronic exposure to 2.9 mT, 40 Hz magnetic field reduces melatonin concentrations in humans" J Pineal Res 25(4): 240-44, 1998.

¹²⁵ Burch JB, Reif JS, Noonan CW, Ichinose T, and Bachand AM. "Melatonin metabolite excretion among cellular telephone users" Int J Radiat Biol 78(11): 1029-36, 2002.

¹²⁶ Lewis AJ, Kerenyl NA, and Feuer G. "Neuropharmacology of Pineal Secretion" Drug Metab Drug Interact 8(3-4): 247-312, 1990.

¹²⁷ Sandyk R and Kay SR. "Abnormal EEG and Calcification of the Pineal Gland in Schizophrenia" Intern J Neurosci 67: 107-11, 1992.

¹²⁸ Bickler SW. "Non-communicable diseases: is their emergence in industrialized societies related to changes in neuroendocrine function?" Med Hypotheses 54(5): 825-8, 2000.

¹²⁹ Weisman AG. "Understanding Cross-Cultural Prognostic Variability for Schizophrenia" Cultural Diversity and Mental Health 3(1): 23-35, 1997.

¹³⁰ Ben-Shachar D. "Mitochondrial dysfunction in schizophrenia: a possible linkage to dopamine" J Neurochem 83: 1241-51, 2002.

¹³¹ Cavelier L, Jazin EE, Eriksson I, Prince J, Bave U, Oreland L, and Gyllesten U. "Decreased cytochrome-c oxidase activity and lack of age-related accumulation of mitochondrial DNA deletions in the brains of schizophrenics" Genomics 29(1): 217-24, 1995.

¹³² Mulcrone J, Whatley SA, Ferrier IN, and Marchbanks RM. "A study of altered gene expression in frontal cortex from schizophrenic patients using differential screening" Schizophr Res 14(3): 203-13, 1995.

¹³³ Dwivedi RS, Dwivedi U, and Chiang B. "Low Intensity Microwave Radiation Effects on the Ultrastructure of Chang Liver Cells" Exp Cell Res 180: 253-65, 1989.

¹³⁴ Sanders AP, Schaefer DJ, and Joines WT. "Microwave Effects on Energy Metabolism of Rat Brain" Bioelectromagnetics 1: 171-81, 1980.

¹³⁵ Sanders AP, Joines WT, and Allis JW. "Effects of Continuous-Wave, Pulsed, and Sinusoidal-Amplitude-Modulated Microwaves on Brain Energy Metabolism" Bioelectromagnetics 6: 89-97, 1985.

McClure RJ, Keshavan MS, and Pettegrew JW. "Chemical and Physiologic Brain Imaging in Schizophrenia" Psychiatr Clin N Am 21(1): 93-122, 1998.

¹³⁶ Baranski S and Edelwejn Z. "Experimental Morphologic and Electroencephalographic Studies of Microwave Effects on the Nervous System" Ann N Y Acad Sci 247: 109-16, 1975.

¹³⁷ Rothermundt M, Arolt V, and Bayer TA. "Review of Immunological and Immunopathological Findings in Schizophrenia" Brain Behav Immunity 15: 319-39, 2001.

¹³⁸ Salford LG, Brun A, Sturesson K, Eberhardt JL, and Persson BR. "Permeability of the blood-brain barrier by 915 MHz electromagnetic radiation, continuous wave and modulated at 8, 16, 50, and 200 Hz" Microsc Res Tech 27(6): 535-42, 1994.

¹³⁹ Oscar KJ and Hawkins TD. "Microwave alteration of the blood-brain barrier system of rats" Brain Res 126(2): 281-93, 1977.

¹⁴⁰ Frey AH, Feld SR, and Frey B. "Neural function and behavior: Defining the relationship in Biological Effects of Nonionizing Radiation" Ann NY Acad Sci 247: 433-38, 1975.

¹⁴¹ Vinogradov GI, Andrienko IG, and Naumenko GM. "[The phenomenon of adaptive immunity in exposure to nonionizing microwave radiation]" Radiobiologiia 31(5): 718-21, 1991.

¹⁴² Neubauer C, Phelan AM, Kues H, and Lange DG. "Microwave irradiation of rats at 2.45 GHz activates pinocytoticlike uptake of tracer by capillary endothelial cells of cerebral cortex" Bioelectromagnetics 11(4): 261-8, 1990.

¹⁴³ Albert EN and Kerns JM. "Reversible microwave effects on the blood-brain barrier" Brain Res 230(1-2): 153-64, 1981.

¹⁴⁴ Leszczynski D, Joenvaara S, Reivinen J, and Kuokka R. "Non-thermal activation of the hsp27/p38MARK stress pathway by mobile phone radiation in human endothelial cells: Molecular mechanism for cancer- and blood-brain barrier-related effects" Differentiation 70: 120-29, 2002.

¹⁴⁵ Frey AH. "Headaches from Cellular Telephones: Are They Real and What Are the Implications" Environ Health Perspect 106(3): 101-3, 1998.

¹⁴⁶ Williams WM, Lu ST, Del Cerro M, and Michaelson SM. "Effect of 2450 MHz microwave energy on the bloodbrain barrier to hydrophilic molecules. D. Brain temperature and blood-brain barrier permeability to hydrophilic tracers" Brain Res 319(2): 191-212, 1984.

¹⁴⁷ Williams WM, Platner J, and Michelson SM. "Effect of 2450 MHz microwave energy on the blood-brain barrier to hydrophilic molecules. C. Effect on the permeability to [14C]sucrose" Brain Res 319(2): 183-90, 1984.

¹⁴⁸ Moriyama E, Saleman M, and Broadwell RD. "Blood-brain barrier alteration after microwave-induced hyperthermia is purely a thermal effect: I. Temperature and Power Measurements" Surg Neurol 35(3): 177-82, 1991.

¹⁴⁹ Gaughran F. "Immunity and Schizophrenia, Autoimmunity, Cytokines, and Immune Responses" Int Rev Neruobiol 52: 275-302, 2002.

¹⁵⁰ Ganguli R, Brar JS, and Rabin BS. "Immune Abnormalities in Schizophrenia Evidence for the Autoimmune Hypothesis" Harvard Rev Psychiatry 2(2): 70-83, 1994.

¹⁵¹ Ganguli R, Rabin BS, Kelly RH, Lyte M, and Ragu U. "Clinical and laboratory evidence of autoimmunity in acute schizophrenia" Ann N Y Acad Sci 496: 676-85, 1987.

¹⁵² Wright P, Sham PC, Gilvarry CM, Jones PB, Cannon M, Sharma T, and Murray RM. "Autoimmune diseases in the pedigrees of schizophrenic and control subjects" Schizophr Res 20(3): 261-7, 1996.

¹⁵³ Grigur'ev VV, Ogurtsov RP, and Zubzhitskii IuN. "[Immunomorphologic changes in the testes upon exposure to a microwave electromagnetic field]" Arkh Anat Gistol Embriol 80(2): 69-75, 1981.

¹⁵⁴ Vinogradov GI and Naumenko GM. "[Experimental modeling of autoimmune reactions as affected by nonionizing radiation]" Radiobiologiia 26(5): 705-8, 1986.

¹⁵⁵ Vinogradov GI, Batanov GV, Naumenko GM, Levin AD, and Trifonov SI. "[Effect of nonionizing microwave radiation on autoimmune reactions and antigenic structure of serum proteins]" Radiobiologiia 25(6): 840-3, 1985.

¹⁵⁶ Rothermundt M, Arolt V, and Bayer TA. "Review of Immunological Findings in Schizophrenia" Brain Behav Immunity 18: 319-39, 2001.

¹⁵⁷ Cossarizza A, Angioni S, Petraglia F, Genazzani AR, Monti D, Capri M, Bersani F, Cadossi R, and Franceschi C. "Exposure to Low Frequency Pulsed Electromagnetic Fields Increases Interleukin-1 and Interleukin-6 Production by Human Peripheral Blood Mononuclear Cells" Exp Cell Res 204: 385-7, 1993. ¹⁵⁸ Fresenko EE, Makar VR, Novoselova EG, and Sadovnikov VB. "Microwaves and cellular immunity I. Effect of whole body microwave irradiation on tumor necrosis factor production in mouse cells" Bioelectrochem Bioenerg 49: 29-35, 1999.

¹⁵⁹ Novoselova EG, Fesenko EE, Makar VR, and Sadovnikov VB. "Microwaves and cellular immunity II. Immunostimulating effects of microwaves and naturally occurring antioxidant nutrients" Bioelectrochem Bioenerg 49: 37-41, 1999.

¹⁶⁰ Novoselova ET and Fesenko EE. "[Stimulation of production of tumor necrosis factory by murine macrophages when exposed in vivo and in vitro to weak electromagnetic waves in the centimeter range" Biofizika 43(6): 1132-3, 1998.

¹⁶¹ Novoselova EG, Ogai VB, Sorokina OV, Novikow VV, and Fesenko EE. "[Effect of centimeter microwaves and the combined magnetic field on the tumor necrosis factor production in cells of mice with experimental tumors]" Biofizika 46(1): 131-5, 2001.

¹⁶² Glushkova OV, Novoselova EG, Sinotova OA, and Vrublevskaia VV. "[Immunomodulation effect of electromagnetic waves on production of tumor necrosis factor in mice with various rates of neoplastic growth]" Biofizika 47(2): 376-81, 2002.

¹⁶³ Wiktor-Jedrzejczak W, Ahmed A, Sell KW, Czerski P, and Leach WM. "Microwaves Induce an Increase in the Frequency of Complement Receptor-bearing Lymphoid Spleen Cells in Mice" J Immunol 118(4): 1499-1502, 1977. ¹⁶⁴ Smialowicz RJ, Brugnolotti PL, and Riddle MM. "Complement Receptor Positive Spleen Cells in Microwave (2450-MHz)-Irradiated Mice" J Microwave Power 16(1): 73-77, 1981.

¹⁶⁵ Wiktor-Jedrzejczak W, Ahmed A, Czerski P, Leach WM, and Sell KW. "Effect of Microwaves (2450-MHz) on the Immune System in Mice: Studies of Nucleic Acid and Protein Synthesis" Bioelectromagnetics 1: 161-70, 1980.

¹⁶⁶ Schlagel CJ, Sulek K, Ho HS, Leach WM, Ahmed A, and Woody JN. "Biological Effects of Microwave Exposure. II. Studies on the Mechanisms Controlling Susceptibility to Microwave-Induced Increases in Complement Receptor-Positive Spleen Cells" Bioelectromagnetics 1: 405-14, 1980.

¹⁶⁷ Schlagel CJ and Ahmed A. "Evidence for Genetic Control of Microwave-Induced Augmentation of Complement Receptor-Bearing B Lymphocytes" J Immunol 129(4): 1530-33, 1982.

¹⁶⁸ Wiktor-Jedrzejczak W, Schlagel CJ, Ahmed A, Leach WM, and Woody JN. "Possible Humoral Mechanism of 2450-MHz Microwave-induced Increase in Complement Receptor Positive Cells" Bioelectromagnetics 2: 81-84, 1981.

¹⁶⁹ Czerska EM, Elson EC, Davis CC, Swicord ML, and Czerski P. "Effects of Continuous and Pulsed 2450-MHz Radiation on Spontaneous Lymphoblastoid Transformation of Human Lymphocytes In Vitro" Bioelectromagnetics 13: 247-59, 1992.

¹⁷⁰ Shenton ME, Dickey CC, Frumin M, and McCarley RW. "A review of MRI findings in schizophrenia" Schizophr Res 49: 1-52, 2001.

¹⁷¹ Heckers S. "Neuroimaging studies of the hippocampus in schizophrenia" Hippocampus 11(5): 520-8, 2001.

Harrison PJ. "The neuropathology of schizophrenia: A critical review of the data and their interpretation" Brain 122(4): 593-624, 1999.

¹⁷² Albert EN and DeSantis M. "Do Microwaves Alter Nervous System Structure" Ann N Y Acad Sci 247: 87-108, 1975.

Kinney DK, Yurgelun-Todd DA, and Woods BT. "Neurologic signs of cerebellar and cortical sensory dysfunction in schizophrenics and their relatives" Schizophr Res 35: 99-104, 1999.

Martin P and Albers M. "Cerebellum and Schizophrenia: A Selective Review" Schizophr Bull 21(2): 241-50, 1995.

¹⁷³ Albert EN, Sherif MF, Papadopoulos NJ, Slaby FJ, and Monahan J. "Effect of Nonionizing Radiation on the Purkinje Cells of the Rat Cerebellum" Bioelectromagnetics 2: 247-57, 1981.

¹⁷⁴ Inouye M, Galvin MJ, and McRee DI. "Effects of 2.45 GHz microwave radiation on the development of Japanese quail cerebellum" Teratology 25(1): 115-21, 1982.

¹⁷⁵ Albert EN and Sherif M. "Morphological changes in cerebellum of neonatal rats exposed to 2.45 GHz microwaves" Prog Clin Biol Res 257: 135-51, 1988.

Lai H. "Neurological Effects of Radiofrequency Electromagnetic Radiation" In: Lin JC (ed.) <u>Advances in</u> <u>Electromagnetic Fields in Living Systems</u> vol 1, Plenum, N Y & London, p 27-80, 1994.

¹⁷⁶ Baranski S. "Histological and Histochemical Effect of Microwave Irradiation on the Central Nervous System of Rabbits and Guinea Pigs" Am J Phys Med 51: 182-91, 1972.

¹⁷⁷ Peinnequin A, Piriou A, Mathieu J, Dabouis V, Sebbah C, Malabiau R, and Debouzy JC. "Non-thermal effects of continuous 2.45 GHz microwaves on Fas-induced apoptosis in human Jurkat T-cell line" Bioelectrochemistry 51(2): 157-61, 2000.

¹⁷⁸ Weiss AP and Heckers S. "Neuroimaging of hallucinations: a review of the literature" Psychiatry Res: Neuroimaging Section 92: 61-74, 1999.

Wilson BS, Zook, JM, Joines WT, and Casseday JH. "Alterations in Activity at Auditory Nuclei of the Rat Induced by Exposure to Microwave Radiation: Autoradiographic Evidence Using [14–C]2Deoxy-D-Glucose" Brain Res 187:

291-306, 1980.

¹⁷⁹ Warwick R and Williams PL (eds.). <u>Gray's Anatomy</u> 35th British Ed, W. B. Saunders Co., 1973.

¹⁸⁰ Oscar KJ, Gruenau SP, Folker MT, and Rapoport SI. "Local cerebral blood flow after microwave exposure" Brain Res 204: 220-25, 1981.

¹⁸¹ Huber R, Treyer V, Borbely AA, Schuderer J, Gottselig JM, Landolt H-P, Werth E, Berthold T, Kuster N, Buck A, and Achermann P. "Electromagnetic fields, such as those from mobile phones, alter regional cerebral blood flow and sleep and waking EEG" J Sleep Res 11: 289-95, 2002.

¹⁸² Thuroczy G, Kubinyi G, Bodo M, Bakos J, and Szabo LD. "Simultaneous Response of Brain Electrical Activity (EEG) and Cerebral Circulation (REG) to Microwave Exposure in Rats" Rev Environ Health 10(2): 135-48, 1994.

¹⁸³ Owega A, Klingelhofer J, Sabri O, Kunert SO, Albers M, and Sass H. "Cerebral blood flow velocity in acute schizophrenic patients: A transcranial Doppler ultrasonography study" Stroke 29(6): 1149-54, 1998.

¹⁸⁴ Hoyer S and Oesterreich K. "Blood flow and oxidative metabolism of the brain in patients with schizophrenia" Psychiatr Clin (Basel) 8(6): 304-13, 1975.

¹⁸⁵ Ohmoto Y, Fujisawa H, Ishikawa T, Koizumi H, Matsuda T, and Ito H. "Sequential changes in cerebral blood flow, early neuropathological consequences and blood-brain barrier disruption following radiofrequency-induced localized hyperthermia in the rat" Int J Hyperthermia 12(3): 321-34, 1996.

¹⁸⁶ David AS. "Auditory hallucinations: phenomenology, neuropsychology and neuroimaging update" Acta Psychiatr Scand 99(Suppl 395): 95-104, 1999.

¹⁸⁷ Taylor SF. "Cerebral blood flow activation and functional lesions in schizophrenia" Schizophr Res 19: 129-40, 1996.

¹⁸⁸ Goldman-Rakic PS and Seleman LD. "Functional and Anatomical Aspects of Prefrontal Pathology in Schizophrenia" Schizophr Bull 23(3): 437-58, 1997.

¹⁸⁹ Bachsbaum MS and Hazlett EA. "Positron Emission Tomography Studies of Abnormal Glucose Metabolism in Schizophrenia" Schizophr Bull 24(3): 343-64, 1998.

¹⁹⁰ Engelien A, Stern E, and Silbersweig D. "Functional Neuroimaging of Human Central Auditory Processing in Normal Subjects and Patients with Neurological and Neuropsychiatric Disorders" J Clin Exp Neuropsychology 23(1): 94-120, 2001.

¹⁹¹ Mohr B, Pulvermuller F, Cohen R, and Rockstroh B. "Interhemispheric cooperation during word processing: evidence for callosal transfer dysfunction in schizophrenic patients" Schizophr Res 46: 231-39, 2000.

¹⁹² Feinberg I, Thode HC, Chugani HT, and March JD. "Gamma Distribution Model Describes Maturational Curves for Delta Wave Amplitude, Cortical Metabolic Rate and Synaptic Density" J Theor Biol 142: 149-61, 1990.

Kendler KS and Davis KL. "The Genetics and Biochemistry of Paranoid Schizophrenia and Other Paranoid Psychoses" Schizophr Bull 7(4): 689-709, 1981.

¹⁹³ Torrey EF. "The Epidemiology of Paranoid Schizophrenia" Schizophr Bull 7(4): 588-93, 1981.

¹⁹⁴ Donald AG, Pressley LC, and Pitts WM Jr. "Changes in the clinical picture of schizophrenia" South Med J 69(11): 1406-9, 1976.

¹⁹⁵ Lung F-W, Tzeng D-S, and Shu B-C. "Ethnic heterogeneity in allele variation in the DRD4 gene in schizophrenia" Schizophr Res 57: 239-45, 2002.

¹⁹⁶ Gorwood P, Leboyer M, Jay M, Payan C, and Feingold J. "Gender and Age at Onset in Schizophrenia: Impact of Family History" Am J Psychiatry 152(2): 208-12, 1993.

¹⁹⁷ Kendler KS and Hays P. "Familial and Sporadic Schizophrenia: A Symptomatic, Prognostic, and EEG Comparison" Am J Psychiatry 139(12): 1557-62, 1982.

¹⁹⁸ Kendell RE, Malcolm DE, and Adams W. "The Problem of Detecting Changes in the Incidence of Schizophrenia" Br J Psychiatry 162: 212-18, 1993.

¹⁹⁹ Munk-Jergensen P. "Decreasing first-admission rates of schizophrenia among males in Denmark from 1970 to 1984" Acta Psychiatr Scand 73: 645-650, 1986.

²⁰⁰ Osby U, Hammar N, Brandt L, Wicks S, Thinsz Z, Ekbom A, and Sparen P. "Time trends in first admissions for schizophrenia and paranoid psychosis in Stockholm County, Sweden" Schizophr Res 47: 247-54, 2001.

²⁰¹ Goldner EM, Hsu L, Waraich P, and Somers JM. "Prevalence and Incidence Studies of Schizophrenic Disorders: A Systematic Review of the Literature" Can J Psychiatry 47(9): 833-43, 2002.

²⁰² Lipman RM, Tripathi BJ, and Tripathi RC. "Cataracts Induced by Microwave and Ionizing Radiation" Surv Ophthal 33(3): 200-10, 1988.

²⁰³ Zaret MM. "Microwave Cataracts" Med Trial Tech Q 19(3): 246-52, 1973.

²⁰⁴ McCarty CA, Wood CA, Fu CL, Livingston PM, Mackersey S, Stanislavsky Y, and Taylor HR. "Schizophrenia, Psychotropic Medication, and Cataract" Ophthalmology 106(4): 683-7, 1999.

²⁰⁵ Ruigomez A, Rodriguez LAG, Dev VJ, Arellano F, and Raniwala J. "Are Schizophrenics or Antipsychotic Drugs a Risk Factor for Cataracts" Epidemol 11(6): 620-3, 2000.

²⁰⁶ Bond WS and Yee GC. "Ocular and cutaneous effects of chronic phenothiazine therapy" Am J Hosp Pharm 37: 74-8, 1980.

²⁰⁷ Aurell E and Tengroth B. "Lenticular and Retinal Changes Secondary to Microwave Exposure" Acta Ophthalmol 51: 764-71, 1973.

²⁰⁸ Paulsson LE, Hamnerius Y, Hansson HA, and Sjostrand J. "Retinal Damage Experimentally Induced by Microwave Radiation at 55 mW/cm²" Acta Ophthmol 57: 183-97, 1979.

²⁰⁹ Reilly SA and Fenton JM. "Thioridazine for schizophrenia" Cochrane Database Syst Rev2000:(3): CD001944.

²¹⁰ Boet DJ. "Phenothiazine Retinopathy" Ophthalmologica Additamentum ad vol. 158: 574-82, 1969.

²¹¹ Steneck NH. <u>The Microwave Debate</u> MIT Press, Cambridge, Mass, London Eng, 1984.

²¹² O'Connor ME. "Psychological Studies in Nonionizing Electromagnetic Energy Research" J Gen Psychol 120(1): 35-47, 1993.

²¹³ Becker RO. <u>Cross Currents</u> Jeremy P. Tarcher, Inc, Los Angeles, St Martin's Press, p 297-304 & p 303-4, 1990.

²¹⁴ Galeev AL. "The Effects of Microwave Radiation from Mobile Telephones on Humans and Animals" Neurosci Behav Physiol 30(2): 187-94, 2000.

²¹⁵ Levallois P. "Hypersensitivity of Human Subjects to Environmental Electric and Magnetic Field Exposure: A Review of the Literature" Environ Health Perspect 110(Supp. 4, Aug.): 613-18, 2002.

²¹⁶ Grundler W, Kaiser F, Keilmann F, and Walleczek J. "Mechanisms of Electromagnetic Interaction with Cellular Systems" Naturwissenschaften 79: 551-9, 1992.

²¹⁷ Richardson-Andrews RC. "Sunspots and the Recency Theory of Schizophrenia" Med Hypotheses 44: 16-19, 1995.

²¹⁸ Lieberman JA. "Is Schizophrenia a Neruodegenerative Disorder? A Clinical and Neurobiological Perspective" Biol Psychiatry 46: 729-39, 1999.

²¹⁹ Knoll JL, Garver DL, Ramberg JE, Kingsbury SJ, Croissant D, and McDermott B. "Heterogeneity of the Psychoses: Is There a Neurodegenerative Psychosis?" Schizophrenia Bull 24(3): 365-79, 1998.

²²⁰ Ahlbom A. "Neurodegenerative Diseases, Suicide and Depressive Symptoms in Relation to EMF" Bioelectromagnetics 5: S132-S143, 2001.

²²¹ Department of the Army, USAF Scientific Advisory Board. "New World Vistas: air and space for the 21st century" 14 vol. (Ancillary Volume) p 89-90, 1996. Also at <u>http://www.pjproject.org/usaf.html</u>

²²² EMF Services. "EMF Shielding & Alternatives" at <u>http://www.emfservices.com/emf-shielding.htm</u>